

EGU2020-1506

<https://doi.org/10.5194/egusphere-egu2020-1506>

EGU General Assembly 2020

© Author(s) 2020. This work is distributed under the Creative Commons Attribution 4.0 License.



Characterising diurnal- and synoptic-timescale changes in urban air quality using Radon-222

Scott Chambers¹, Dafina Kikaj², Agnieszka Podstawczyńska³, Jagoda Crawford¹, and Alastair Williams¹

¹Environmental Research, ANSTO, Lucas Heights, Australia

²Dept. Environmental Sciences, Jožef Stefan Institute, Ljubljana, Slovenia

³Dept. Meteorology and Climatology, University of Łódź, Łódź, Poland

Urban air quality is strongly influenced by the atmosphere's ability to disperse primary emissions and opportunities for secondary pollution formation. In mid- to high-latitude regions that experience enduring winter snow cover or soil freezing, regional subsidence and stagnation associated with persistent anti-cyclonic conditions such as the "Siberian High" can lead to "cold pool" or "persistent inversion" events. These events can result in life-threatening pollution episodes that last for weeks. While often associated with complex topography [1,2], persistent inversion events can also influence the air quality of urban centres in flat, inland regions [3]. This presentation will describe a recently-developed radon-based technique for identifying and characterising synoptic-timescale persistent inversion events, which is proving to be a simple and economical alternative to contemporary meteorological approaches that require regular sonde profiles [1]. Furthermore, key assumptions of the radon-based technique to characterise diurnal-timescale changes in the atmospheric mixing state described by Chambers et al. [4] are violated during persistent inversion conditions. Here we demonstrate how atmospheric class-typing, through successive application of radon-based techniques for identifying synoptic- and diurnal-timescale changes in the atmospheric mixing state, improves understanding of atmospheric controls on urban air quality in non-summer months across the full diurnal cycle. This knowledge translates directly to statistically-robust techniques for assessing public exposure to pollution, and for evaluating the efficacy of pollution mitigation measures. Lastly, we show how atmospheric class-typing can be used to enhance the evaluation of chemical transport models [5].

[1] Baasandorj, M., et al. *Environ. Sci. Technol.*, 51, 5941–5950, <https://doi.org/10.1021/acs.est.6b06603>, 2017.

[2] Kikaj, D., et al. *Atmos. Meas. Tech.*, 12, 4455–4477, <https://doi.org/10.5194/amt-12-4455-2019>, 2019.

[3] Chambers, SD and A Podstawczyńska. *Atmos. Environ.*, 219, 117040, <https://doi.org/10.1016/j.atmosenv.2019.117040>, 2019.

[4] Chambers, S.D., et al. *J. Geophys. Res. Atmos.* 124, 770–788,

<https://doi.org/10.1029/2018JD029507>, 2019.

[5] Chambers, S.D., et al. *Atmosphere* 10 (1), 25, doi:10.3390/atmos10010025, 2019.